

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
23 February 2006 (23.02.2006)

(10) International Publication Number
WO 2006/018807 A1

(51) International Patent Classification:
C07D 501/00 (2006.01) A61P 31/04 (2006.01)
A61K 31/546 (2006.01)

(74) Common Representative: RANBAXY LABORATO-
RIES LIMITED; c/o DESHMUKH, Jay R., 600 College
Road East, Suite 2100, Princeton, NJ 08540 (US).

(21) International Application Number:
PCT/IB2005/052691

(22) International Filing Date: 15 August 2005 (15.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1508/DEL/2004 16 August 2004 (16.08.2004) IN
1646/DEL/2004 31 August 2004 (31.08.2004) IN
434/DEL/2005 28 February 2005 (28.02.2005) IN

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): RAN-
BAXY LABORATORIES LIMITED [IN/IN]; Plot No.
90, Sector - 32, Gurgaon, Haryana 122 001 (IN).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): MAHENDRU,
Manu [IN/IN]; 5733, Ground Floor, Sector - 38 West,
Chandigarh, Chandigarh 160014 (IN). ARYAN, Ram,
Chander [IN/IN]; 1066, Sector - A, Pocket - A, Vasant
Kunj, New Delhi, Delhi 110070 (IN). KUMAR, Satish
[IN/IN] (IN). PANDYA, Bhargav [IN/IN]; 7, Vallabh
Apartments, Maniyasha, Maninagar (East), Ahmed-
abad, Gujarat 380008 (IN). DUGGAL, Sanjam [IN/IN]
(IN). GADE, Sanjay [IN/IN] (IN). KUMAR, Yatendra
[IN/IN]; U-26/5, Phase - III/IDLF Qutab Enclave, Gurgaon,
Haryana 122001 (IN).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CRYSTALLINE FORMS OF CEFIDINIR



(57) Abstract: The invention relates to processes for the preparation of crystalline polymorphic forms of cefdinir of formula (I). More particularly, it relates to the preparation of crystalline polymorphic forms of cefdinir designated as Forms B and C. The invention also relates to pharmaceutical compositions that include the polymorphic forms B and C, and the use of the compositions for treating bacterial infections.

WO 2006/018807 A1

Description

CRYSTALLINE FORMS OF CEFDINIR

[1] Field of the Invention


[2] The field of the invention relates to processes for the preparation of crystalline polymorphic forms of cefdinir. More particularly, it relates to the preparation of crystalline polymorphic forms of cefdinir designated as Forms B and C. The invention also relates to pharmaceutical compositions that include the polymorphic Forms B and C, and the use of the compositions for treating bacterial infections.

[3] Background of the Invention

[4] Chemically, cefdinir is $[(-)6R,7R]$ -

7-((Z)-2-(2-amino-4-thiazolyl)-2-hydroxyiminoacetamido)-3-vinyl-8-oxo-5-thia-1-aza bicyclo[4.2.0]oct-2-ene-2-carboxylic acid having the structural Formula I. It is a third-generation semi-synthetic cephalosporin for oral use, characterized by broad antibacterial spectrum against gram-positive and gram-negative bacteria. In particular, the cefdinir shows excellent antibacterial activity against *Streptococci* and *Staphylococci*.

[5]



[6] **FORMULA I**

[7] U.S. Patent No. 4,559,334 discloses a process for the preparation of cefdinir in amorphous form by lyophilization. The amorphous form so produced is highly hygroscopic and therefore very difficult to formulate.

[8] U.S. Patent No. 4,935,507 discloses a crystalline Form A of cefdinir having a specific X-Ray diffraction pattern as shown in Figure 1.

[9] U.S. Application No. 20030204082 discloses a crystalline form of cefdinir (hereinafter designated as Form R) having a specific X-Ray diffraction pattern as shown in Figure 2.

[10] U.S. Application No. 20040242556 discloses a crystalline form of cefdinir designated as Form B having a specific X-Ray diffraction pattern. Form B is prepared from crystalline Form A by first forming a trifluoroacetic acid salt followed by basification with ammonia.

- [11] U.S. Patent No. 6,350,869 describes a process for the preparation of the crystalline dicyclohexylamine salt of cefdinir and monohydrate form of cefdinir.
- [12] International (PCT) Publication No. WO 04/46154 describes the amorphous monohydrate of cefdinir and a process for preparation thereof.
- [13] Several processes have been reported for the preparation of crystalline salts of cefdinir which are either useful as intermediates in preparation of cefdinir or can be used as broad spectrum antimicrobials, for example, in International (PCT) Publication Nos. WO 02/98884; 04/16623; and 04/56835.
- [14] International (PCT) Publication No. WO 03/50124 describes crystalline cefdinir potassium monohydrate and process for preparation thereof. The application further details utilization of the potassium salt of cefdinir in monohydrate form as potential antimicrobial agent.
- [15] The details of one or more embodiments of the inventions are set forth in the description below. Other features, objects and advantages of the inventions will be apparent from the description, figures, and claims.
- [16] Description of the Drawings
- [17] Figure 1 is an X-ray powder diffraction pattern of Form A of cefdinir.
- [18] Figure 2 is an X-ray powder diffraction pattern of crystalline cefdinir obtained as per U.S. Application No. 20030204082.
- [19] Figure 3 is an X-ray powder diffraction pattern of crystalline Form B of cefdinir.
- [20] Figure 4 is a differential scanning calorimetry thermogram of crystalline Form B of cefdinir.
- [21] Figure 5 is an infrared spectrum of crystalline Form B of cefdinir.
- [22] Figure 6 is an X-ray powder diffraction pattern of Form C of cefdinir.
- [23] Figure 7 is a differential scanning calorimetry thermogram of polymorphic Form C of cefdinir.
- [24] Figure 8 is an infrared spectrum of crystalline Form C of cefdinir.
- [25] Summary of the invention
- [26] In one general aspect there is provided a crystalline Form B of cefdinir.
- [27] The Form B of cefdinir may have the X-ray diffraction pattern of Figure 3, differential scanning calorimetry thermogram of Figure 4, and infrared spectrum of Figure 5.
- [28] In another general aspect there is provided a pharmaceutical composition that includes a therapeutically effective amount of the crystalline Form B of cefdinir; and one or more pharmaceutically acceptable carriers, excipients or diluents.
- [29] In one general aspect there is provided a crystalline Form C of cefdinir.
- [30] The Form C of cefdinir may have the X-ray diffraction pattern of Figure 6, differential scanning calorimetry thermogram of Figure 7, and infrared spectrum of Figure

8.

- [31] In another general aspect there is provided a pharmaceutical composition that includes a therapeutically effective amount of the crystalline Form C of cefdinir; and one or more pharmaceutically acceptable carriers, excipients or diluents.
- [32] In another general aspect there is provided a process for the preparation of crystalline form of cefdinir. The process includes converting cefdinir or a base addition salt thereof to cefdinir acid addition salt with an acid at a temperature of about 10°C or less; basifying the acid addition salt with a base; and isolating crystalline cefdinir.
- [33] The process may include further drying of the product obtained.
- [34] The process may produce Form B or Form C of crystalline cefdinir. It may produce the hydrated form of crystalline cefdinir.
- [35] In another general aspect there is provided a pharmaceutical composition that includes a therapeutically effective amount of one or both of the crystalline Form B and the crystalline Form C of cefdinir, and one or more pharmaceutically acceptable carriers, excipients or diluents.
- [36] Embodiments of the pharmaceutical compositions may include one or more of the following features or those described above. For example, the cefdinir may be either substantially pure crystalline Form B cefdinir or substantially pure crystalline Form C cefdinir.
- [37] In another general aspect there is provided a method of treating bacterial infections in a warm-blooded animal. The method includes providing a pharmaceutical composition to the warm-blooded animal, the pharmaceutical composition comprising one or both of the crystalline Form B and the crystalline Form C of cefdinir.
- [38] Embodiments of the method may include one or more of the following features or those described above. For example, the cefdinir may be either substantially pure crystalline Form B cefdinir or substantially pure crystalline Form C cefdinir.
- [39] Detailed Description of the Invention
- [40] The inventors have developed a novel process for preparation of crystalline polymorphic forms of cefdinir. The present inventors also have surprisingly found new polymorphic forms of cefdinir, which are characteristically different from the existing crystalline forms of cefdinir. The new polymorphic forms are designated as Forms B and C of cefdinir. All these forms are found to be stable under normal and accelerated storage conditions.
- [41] Hereinafter the term 'crystalline cefdinir' refers to any crystalline polymorphic form of cefdinir. The term 'Form B' of cefdinir refers to crystalline cefdinir having X-Ray Powdered Diffraction (XRPD) pattern as depicted in Figure 3. The term 'Form C' of cefdinir refers to crystalline cefdinir having X-Ray Powdered Diffraction (XRPD) pattern as depicted in Figure 5.

- [42] A first aspect of the present invention provides a process for the preparation of crystalline cefdinir wherein the process includes the steps of:
- [43] (a) converting cefdinir to cefdinir acid addition salt at a temperature of 10°C or less;
- [44] (b) optionally isolating the acid addition salt obtained in step a);
- [45] (c) basifying the acid addition salt of cefdinir with a base; and
- [46] (d) isolating crystalline cefdinir from the reaction mixture thereof.
- [47] Cefdinir is suspended in an organic solvent or water at a temperature of 10°C or less. The resultant mixture is acidified with a suitable acid. The resultant mass is stirred and the precipitated solids are filtered, washed with water. The product obtained is dried to yield cefdinir acid addition salt.
- [48] The organic solvent can be selected from one or more of esters, lower alkanols, ethers, ketones, polar aprotic solvents, halogenated hydrocarbons or mixtures thereof.
- [49] The acid used is one or more inorganic or organic acids selected from one or more of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, trifluoroacetic acid (TFA), p-toluenesulphonic acid (pTSA), maleic acid, acetic acid, succinic acid and the like.
- [50] The acid addition salt is basified with a base to a pH of about 1 to about 3, stirred and the precipitated solids are filtered, washed with water. The product obtained is dried to yield crystalline cefdinir. Forms B and C of cefdinir can be easily prepared by following the processes of the present invention.
- [51] The base used can be selected from one or more inorganic or organic bases such as alkali and alkaline earth metal hydroxide or alkoxide, ammonia, triethylamine, dicyclohexylamine amine or diisopropylamine and the like.
- [52] A second aspect of the present invention provides a process for the preparation of crystalline cefdinir wherein the process includes the steps of:
- [53] (a) converting a base addition salt of cefdinir to cefdinir acid addition salt;
- [54] (b) optionally isolating the acid addition salt of cefdinir,
- [55] (c) basifying the acid addition salt of cefdinir with a base; and
- [56] (d) isolating crystalline cefdinir from the reaction mixture thereof.
- [57] Base addition salt of cefdinir is a salt of cefdinir with inorganic or organic base. Several such base addition salts are known to skilled artisans. The base addition salt of cefdinir is suspended in an organic solvent and acidified using a suitable acid to adjust the pH of this mixture to between about 1 to about 3. The resultant mass is stirred and the mixture is suitably worked up to provide an acid addition salt of cefdinir. The acid addition salt is then converted to crystalline form of cefdinir by basification as described in the first aspect of the invention.
- [58] The organic solvent optionally contains water and can be selected from one or more of esters, lower alkanols, ethers, ketones, polar aprotic solvents, halogenated hy-

drocarbons or mixtures thereof.

[59] The suitable acid, which is used for preparation of the acid addition salt, and the base, which is used for basifying the acid addition salt, have already been described above.

[60] A third aspect of the present invention provides a process for the preparation of crystalline cefdinir wherein the process includes the steps of:

[61] (a) acidifying an isolated base addition salt of cefdinir at a temperature of 10°C or less; and

[62] (b) isolating crystalline cefdinir from the reaction mixture thereof.

[63] A suspension or solution of an isolated base addition salt of cefdinir is acidified using a suitable acid described earlier at a temperature of less than 10°C. The separated crystalline product is filtered, washed with water and dried to get crystalline cefdinir.

[64] Forms B and C of cefdinir can be easily prepared by following the processes described in the first, second or third aspects of the present invention.

[65] A fourth aspect of the present invention relates to a process for preparation of crystalline Form B of cefdinir wherein the process includes the steps of:

[66] (a) dissolving cefdinir or salt thereof in water at pH of about 5.5 to 8;

[67] (b) optionally adding an organic solvent in the solution of step a);

[68] (c) adjusting the pH of solution obtained in step a) or b) between 1.0 and 3.0 with an acid; and

[69] (d) isolating crystalline Form B of cefdinir from the reaction mixture obtained thereof.

[70] Cefdinir or salt thereof to be used as a starting material can be prepared by conventional methods reported in the literature, including U.S. Patent Nos. 4,559,334; 4,935,507; 4,870,168; 6,350,869 and 6,093,814; PCT Patent Application Nos. WO 98/45299; WO 99/55710; WO 02/98884; WO 03/91261; WO 04/16623; WO 04/35800; WO 04/56835; WO 04/58695 and WO 04/46154.

[71] Cefdinir or salt thereof is first suspended in water and the resultant mixture is cooled to about -20 to 25°C. Preferably the temperature is adjusted in the range of -10 to 20°C. To this mixture a suitable base is added in order to adjust the pH of the reaction mass in the range of 5.5 to 8. The bases are common bases known to persons of ordinary skills in the art and include one or more of alkali or alkaline earth metal hydroxide, alkali or alkaline earth metal carbonate, alkali or alkaline earth metal bicarbonate, alkali or alkaline earth metal alkoxide, alkali or alkaline earth metal hydride or an amine. The amine can be a primary, secondary or tertiary amine having alkyl, aryl, aralkyl, cycalkyl or heterocycle as substituent groups.

[72] A clear solution thus obtained can be further diluted with a suitable organic solvent selected from one or more of water soluble alkanols, ketones, esters, acetonitrile,

tetrahydrofuran, 1,4-dioxane, N,N-dimethylacetamide, N,N-dimethylformamide, N-methylpyrrolidone and dimethylsulphoxide or mixtures thereof.

[73] The pH of the clear solution is then adjusted using an acid in the range of 1.0 to 3.0. The acid can be a mineral acid or an organic acid such as carboxylic or sulphonic acid.

[74] The resultant mass is stirred at 0 to 30°C for 5 to 30 hours and the precipitated solids are filtered and washed with water. The product obtained is dried at room temperature to yield crystalline Form B of cefdinir.

[75] A fifth aspect of the present invention relates to a process for preparation of polymorphic Form C of cefdinir in which the process includes the steps of:

[76] (a) dissolving cefdinir or salt thereof in water at pH of about 5.5 to 8;

[77] (b) optionally adding an organic solvent in the solution of step a);

[78] (c) adjusting the pH of solution obtained in step a) or b) to between 1.0 and 3.5 with an acid; and

[79] (d) isolating the polymorphic Form C of cefdinir from the reaction mixture obtained thereof.

[80] Cefdinir or salt thereof to be used as starting material can be prepared by conventional methods reported in the literature, such as in U.S. Patent Nos. 4,559,33; 4,935,507; 4,870,168; 6,350,869 and 6,093,814; PCT Patent Application Nos. WO 98/45299; WO 99/55710; WO 02/98884; WO 03/91261; WO 04/16623; WO 04/35800; WO 04/56835; WO 04/58695 and WO 04/46154.

[81] Cefdinir or salt thereof is first suspended in water and the resultant mixture is cooled to about -20 to 25°C. Preferably the temperature is adjusted in the range of -10 to 20°C. To this mixture a suitable base is added in order to adjust the pH of the reaction mass in the range of 5.5 to 8. The bases are common bases known to persons of ordinary skills in the art and include one or more of alkali or alkaline earth metal hydroxide, alkali or alkaline earth metal carbonate, alkali or alkaline earth metal bicarbonate, alkali or alkaline earth metal alkoxide, alkali or alkaline earth metal hydride or ammonia or an amine. The amine can be primary, secondary or tertiary amine having alkyl, aryl, aralkyl, cycalkyl or heterocycle as substituent groups.

[82] The clear solution thus obtained can be decolorized using charcoal and further diluted with a suitable organic including one or more solvent selected from alkanols, esters, ketones, acetonitrile, chlorinated hydrocarbons, tetrahydrofuran, N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, 1,4-dioxane and dimethylsulphoxide or mixtures thereof.

[83] The pH of the clear solution is then adjusted using an acid in the range of 1.0 to 3.5. The acid can be mineral acid or an organic acid such as carboxylic or sulphonic acid.

[84] The resultant mass is stirred at 0 to 25°C for sufficient time to precipitate Form C of cefdinir. The separated solids are filtered and washed with water. The product obtained

is dried at room temperature to yield crystalline Form C of cefdinir.

[85] A sixth aspect of the present invention provides crystalline Form B of cefdinir. A seventh aspect of the present invention provides crystalline Form B of cefdinir having a characteristic XRD pattern as depicted in Figure 3.

[86] An eighth aspect of the present invention provides crystalline Form B of cefdinir having characteristic absorption bands at two-theta values of 8.34 ± 0.2 , 10.56 ± 0.2 , 11.8 ± 0.2 , 14.04 ± 0.2 , 15.04 ± 0.2 , 17.92 ± 0.2 , 18.54 ± 0.2 , 18.90 ± 0.2 , 20.52 ± 0.2 , 21.06 ± 0.2 , 21.22 ± 0.2 , 22.08 ± 0.2 , 23.62 ± 0.2 , 25.08 ± 0.2 , 25.72 ± 0.2 , 26.16 ± 0.2 , 27.44 ± 0.2 , 28.24 ± 0.2 , 28.50 ± 0.2 , 30.42 ± 0.2 , 32.00 ± 0.2 , 35.84 ± 0.2 and 38.82 ± 0.2 .

[87] A ninth aspect of the present invention provides crystalline cefdinir having characteristic DSC melting exotherms obtained at about 78-85°C and about 160-170°C. The DSC thermogram of cefdinir is provided as Figure 4.

[88] A tenth aspect of the present invention provides crystalline Form B of cefdinir having a characteristic Fourier Transform Infrared (FTIR) spectrum as depicted in Figure 5.

[89] An eleventh aspect of the present invention provides crystalline cefdinir having a moisture content of about 12 to 14% w/w as measured by Thermal Gravimetric Analysis (TGA) and Karl Fischer analysis.

[90] A twelfth aspect of the present invention provides the crystalline Form C of cefdinir. A thirteenth aspect of the present invention provides crystalline Form C of cefdinir having a characteristic XRD pattern as depicted in Figure 6.

[91] A fourteenth aspect of the present invention provides crystalline Form C of cefdinir having characteristic absorption bands at two-theta values of 9.12 ± 0.2 , 10.72 ± 0.2 , 15.04 ± 0.2 , 17.96 ± 0.2 , 18.66 ± 0.2 , 20.92 ± 0.2 , 21.44 ± 0.2 , 22.32 ± 0.2 , 23.66 ± 0.2 , 24.18 ± 0.2 , 25.72 ± 0.2 , 26.26 ± 0.2 , 27.48 ± 0.2 , 28.30 ± 0.2 , 30.42 ± 0.2 , 32.06 ± 0.2 , 35.76 ± 0.2 , 38.80 ± 0.2 and 39.22 ± 0.2 .

[92] A fifteenth aspect provides crystalline Form C of cefdinir having a differential scanning calorimetric (DSC) thermogram as depicted in Figure 7.

[93] A sixteenth aspect of the present invention provides crystalline cefdinir having a characteristic melting exotherm obtained at about 120-130°C.

[94] A seventeenth aspect of the present invention provides crystalline Form C of cefdinir having characteristic Fourier Transform Infrared (FTIR) spectrum as depicted in Figure 8.

[95] An eighteenth aspect of the present invention provides a pharmaceutical composition comprising crystalline Form B or C of cefdinir along with pharmaceutically acceptable carrier(s) and/or excipient(s). The pharmaceutical compositions include one or more oral dosage forms such as tablets, capsules, liquid orals,

suspensions and the like, as well as topical dosage forms such as creams, lotions, ointments and the like.

[96] A nineteenth aspect of the present invention provides a method of treating bacterial infections comprising administering to a mammal in need thereof a therapeutically effective amount of crystalline Form B or C of cefdinir.

[97] Powder XRD of the samples were determined by using X-Ray Diffractometer, Rigaku Corporation, RU-H3R, Goniometer CN2155A3, X-Ray tube with Cu target anode, Divergence slits 1 0, Receiving slit 0.15mm, Scatter slit 1 , Power: 40 KV, 100 mA, Scanning speed: 2 deg/min step: 0.02 deg, Wave length: 1.5406 A

[98] FT-IR of the samples were determined by using a Perkin Elmer instrument, 16 PC, SCAN: 16 scans, 4.0 cm-1, according to the USP 25, general test methods page 1920, infrared absorption spectrum by potassium bromide pellet method.

[99] DSC thermograms were recorded using DSC821 e, Mettler Toledo, Sample weight: 3-5 mg, Temperature range: 50-350°C, Heating rate: 10°C/min, Nitrogen 80.0 mL/min, Number of holes in the crucible: 1.

[100] While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.

[101] Example 1: Preparation of crystalline Form B of cefdinir

[102] Cefdinir potassium salt (10 g) was suspended in de-mineralized water (300 ml) and cooled to 0-5°C. Methanol (60 ml) was added to this mixture followed by cooling again to 0-5°C. Dilute hydrochloric acid cooled to 0-5°C was added in one lot till pH of the solution was 2.2-2.3. The reaction mixture was stirred for 5 hours at 12-14°C. The separated solids were filtered, washed with de-mineralized water and dried to obtain the title compound. The yield was eight grams. The moisture contents was 13.69% w/w.

[103] Example 2: Preparation of cefdinir hydrochloride

[104] Cefdinir (80 g) was suspended in a mixture of denatured spirit (80 ml) and ethyl acetate (160 ml) at 0-5°C. Concentrated hydrochloric acid (17 ml) was added drop wise at 0-5°C to this mixture to get a clear solution. This clear acidified solution was added drop wise to ethyl acetate (3200 ml) at 0-5°C and the resultant mass was stirred vigorously for 1-2 hours. The precipitated solid was filtered, washed with ethyl acetate (500 ml), and dried to give the title compound. The yield was 96 grams.

[105] Example 3: Preparation of crystalline Form C of cefdinir

[106] Cefdinir hydrochloride (12 g) was dissolved in a mixture of tetrahydrofuran (30 ml) and water (150 ml). The solution was cooled to 18°C followed by addition of dichloromethane (10 ml). Dilute ammonia solution was added drop wise to the resultant mixture at 18-20°C till pH of 2.2 was reached. The reaction mixture was

stirred for 5 hours and the separated solid was filtered, washed with de-mineralized water (30 ml x 3) and dried to obtain the title compound. The yield was 9.2 grams. The moisture content was 12-14% w/w.

[107] Example 4: Preparation of cefdinir hydrochloride

[108] Cefdinir potassium salt (5 g) was suspended in de-mineralized water (25 ml). Concentrated hydrochloric acid (3.5 ml) was added to this mixture slowly till pH of about 1 was reached. Tetrahydrofuran (50 ml) was added to this acidified mixture and stirred for 5 minutes. Sodium chloride (10 g) was added to the reaction mixture followed by stirring. The organic layer was separated and tetrahydrofuran was recovered to yield an oily mass. Ethyl acetate (10 ml) and denatured spirit (5 ml) were added to the oily mass. The resultant mixture was cooled to 0-5°C. To this cooled mixture, concentrated hydrochloric acid (0.85 ml) was slowly added to get a clear solution. This clear acidified solution was added to ethyl acetate (160 ml) at 0-5°C followed by stirring for 1 hour. The separated solid was filtered and dried to obtain the title compound. The yield was 3.5 grams.

[109] Example 5: Preparation of cefdinir para-toluenesulphonic acid salt

[110] Cefdinir (20 g) was suspended in tetrahydrofuran (50 ml) and ethyl acetate (40 ml) at 0-5°C. To this mixture, para-toluenesulphonic acid was added (10.6 g) followed by stirring for 15-20 min to get a clear solution. This clear acidified solution was added drop-wise to previously cooled (0-5°C) ethyl acetate (1 L). The reaction mass was stirred for 1-2 hours at 0-5°C, filtered, and washed with ethyl acetate (100 ml). The separated solid was dried to obtain the title compound. The yield was twenty eight grams.

[111] Example 6: Preparation of crystalline Form C of cefdinir

[112] Cefdinir para-toluene sulphonic acid salt (12 g) was dissolved in a mixture of tetrahydrofuran (30 ml) and water (150 ml) and cooled to 18°C. To this solution, dichloromethane (10 ml) was added. Dilute ammonia was added drop-wise to the resultant mixture till a pH of 2.2 was reached. The reaction mixture was stirred for 5 hrs at 18°C. The separated solid was filtered, washed with de-mineralized water (30 ml x 3), and dried to obtain the title compound. The yield was 9.2 grams and the moisture content was 12-14%.

[113] Example 7: Preparation of cefdinir trifluoroacetic acid salt

[114] Cefdinir (5 g) was suspended in de-mineralized water (200 ml) at room temperature. Trifluoroacetic acid (15 ml) was added drop-wise to this mixture to get a clear solution. This clear acidified solution was stirred for 0.5 hour. The separated solid was filtered, washed with cold water and dried to obtain the title compound. The yield was 4.8 grams.

[115] Example 8: Preparation of crystalline Form C of cefdinir

- [116] Cefdinir trifluoroacetic acid salt (12 g) was dissolved in a mixture of tetrahydrofuran (30 ml) and water (150 ml) and cooled to 18°C. Dichloromethane (10 ml) was added to this solution. Dilute ammonia was added drop-wise to the resultant mixture till a pH of 2.2 was reached. The reaction mixture was stirred for 5 hrs at 18°C. The separated solid was filtered, washed with de-mineralized water (30 ml x 3) and dried to obtain the title compound. The yield was 9.2 grams and the moisture content was 12-14%.
- [117] Example 9: Preparation of cefdinir hydrochloride
- [118] Cefdinir potassium salt (20 g) was suspended in de-mineralized water (100 ml) at room temperature. Denatured spirit (30 ml) and ethyl acetate (100 ml) were added to this mixture. The resultant mixture was cooled to 0-4°C. To the cooled mixture, sodium chloride (32 g) was added followed by stirring for 2-3 minutes. Concentrated hydrochloric acid (8.8 ml, 2.2 M) was added drop wise to the resultant mixture. The organic layer was separated from this acidified mixture. The aqueous layer was washed again with a mixture of denatured spirit (15 ml) and ethyl acetate (50 ml) and the washings added to the previously separated organic layer. The combined organic layers were mixed and concentrated to about half the volume at 0-5°C. The resulting mass was added drop-wise to previously cooled (0-4°C) ethyl acetate (800 ml). The resultant mixture was stirred at 0-4°C for 1-1.5 hours and then filtered, washed with ethyl acetate (50 ml x 2) and dried to obtain the title compound. The yield was nineteen grams.
- [119] Example 10: Preparation of crystalline Form C of cefdinir
- [120] Cefdinir hydrochloride (18 g) was dissolved in a mixture of tetrahydrofuran (45 ml) and de-mineralized water (75 ml). This solution was stirred at room temperature and de-mineralized water (165 ml) was added followed by stirring at 18-19°C. Dilute ammonia solution was added drop-wise at 18-20°C to this mixture till a pH of 2.2-2.3 was reached. This reaction mixture was stirred for 5 hours at 18-19°C, filtered, washed with water (45 ml x 3) and dried to obtain the title compound. The yield was 11.5 grams and the moisture content was 12.84%.
- [121] Example 11: Preparation of crystalline Form B of cefdinir
- [122] Cefdinir potassium salt (10 g) was suspended in water (300 ml) and methanol (60 ml). This mixture was cooled to 0-4°C. To this cooled mixture, dilute hydrochloric acid pre-cooled to 5°C was added till pH of 2.2 was reached at 0-4°C. The reaction mixture was stirred for 5-6 hours at 12-14°C, filtered, washed with de-mineralized water (30 ml x 3) and dried to obtain the title compound. The yield was 8.9 grams and the moisture content was 13.7%.
- [123] Example 12: Preparation of cefdinir hydrochloride
- [124] Crystalline form B cefdinir (7.5 g) was suspended in a mixture of denatured spirit

(4.5 ml) and ethyl acetate (15 ml) at room temperature. This mixture was cooled to 0-5°C and concentrated hydrochloric acid (1.7 ml) was added drop wise at 0-5°C to get a clear solution. This clear acidified solution was added to pre-cooled ethyl acetate (300 ml) at 0-5°C. The resultant mixture was stirred for 1 hour, filtered, washed with ethyl acetate (20 ml x 3) and dried to obtain the title compound. The yield was 6.75 grams.

[125] Example 13: Preparation of crystalline Form B of cefdinir

[126] A stirred suspension of cefdinir (5 g) in water was cooled to about 0°C. The solid was dissolved by adjusting the pH of the mixture to about 5.5 to 8.0. Methanol was added to the clear solution followed by addition of dilute hydrochloric acid and water to adjust the pH at about 1.0 to 2.8 while maintaining the temperature of the reaction mass between 5 to 20°C. The resultant mass was stirred for about 12 to 14 hours and the precipitated solids were filtered and washed. The product was dried at room temperature to yield the crystalline Form B of cefdinir. The yield was 4.9 grams and the moisture content was 12% w/w.

[127] Example 14: Preparation of crystalline Form C of cefdinir

[128] A stirred suspension of cefdinir (5 g) in water was dissolved by adjusting the pH of the mixture to about 5.5 to 8.0. The clear solution was decolorized using activated charcoal (0.5 gm) and the resultant mixture was filtered. A mixture of tetrahydrofuran and dichloromethane (10 to 40 ml) was added to the clear filtrate. The solution so obtained was stirred at 5 to 20°C followed by addition of dilute hydrochloric acid to adjust the pH at about 2.0 to 3.0 while maintaining the temperature of the reaction mass between 5 to 20°C. The resultant mass was stirred for sufficient time to precipitate solids and the precipitated solids were filtered and washed. The product was dried at 40°C to yield the crystalline Form C of cefdinir. The yield was 4.79 grams and the moisture content was 12.0% w/w.

[129] While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention. For example, it is understood that the various polymorphic forms of cefdinir can be incorporated in dosage forms for treating conditions for which cefdinir is useful.

Claims

- [1] Crystalline Form B of cefdinir having the X-ray diffraction pattern of Figure 3.
- [2] The crystalline Form B of cefdinir of claim 1 having characteristic X-ray diffraction peaks at two-theta values of 8.34 ± 0.2 , 10.56 ± 0.2 , 11.8 ± 0.2 , 14.04 ± 0.2 , 15.04 ± 0.2 , 17.92 ± 0.2 , 18.54 ± 0.2 , 18.90 ± 0.2 , 20.52 ± 0.2 , 21.06 ± 0.2 , 21.22 ± 0.2 , 22.08 ± 0.2 , 23.62 ± 0.2 , 25.08 ± 0.2 , 25.72 ± 0.2 , 26.16 ± 0.2 , 27.44 ± 0.2 , 28.24 ± 0.2 , 28.50 ± 0.2 , 30.42 ± 0.2 , 32.00 ± 0.2 , 35.84 ± 0.2 and 38.82 ± 0.2 .
- [3] The crystalline Form B of cefdinir of claim 1 having the differential scanning calorimetry thermogram of Figure 4.
- [4] The crystalline Form B of cefdinir of claim 1 having characteristic differential scanning calorimetric melting exotherms at about 78-85°C and about 160-170°C.
- [5] The crystalline Form B of cefdinir of claim 1 having the infrared spectrum of Figure 5.
- [6] Crystalline cefdinir having a moisture content of about 12 to 14% w/w as measured by Thermal Gravimetric Analysis and Karl Fischer analysis.
- [7] Crystalline Form C of cefdinir having the X-ray diffraction pattern of Figure 6.
- [8] The crystalline Form C of cefdinir of claim 7 having characteristic X-ray diffraction peaks at two-theta values of 9.12 ± 0.2 , 10.72 ± 0.2 , 15.04 ± 0.2 , 17.96 ± 0.2 , 18.66 ± 0.2 , 20.92 ± 0.2 , 21.44 ± 0.2 , 22.32 ± 0.2 , 23.66 ± 0.2 , 24.18 ± 0.2 , 25.72 ± 0.2 , 26.26 ± 0.2 , 27.48 ± 0.2 , 28.30 ± 0.2 , 30.42 ± 0.2 , 32.06 ± 0.2 , 35.76 ± 0.2 , 38.80 ± 0.2 and 39.22 ± 0.2 .
- [9] The crystalline Form C of cefdinir of claim 7 having the differential scanning calorimetric thermogram of Figure 7.
- [10] The crystalline Form C of cefdinir of claim 7 having the characteristic differential scanning calorimetric melting exotherm at about 120-130°C.
- [11] The crystalline Form C of cefdinir of claim 7 having the infrared spectrum of Figure 8.
- [12] A process for the preparation of crystalline cefdinir, the process comprising:
(a) converting cefdinir or a base addition salt thereof to cefdinir acid addition salt with an acid at a temperature of about 10°C or less;
(b) basifying the acid addition salt of cefdinir with a base; and
(c) isolating crystalline cefdinir.
- [13] The process of claim 12, wherein the acid comprises an inorganic or organic acid.
- [14] The process of claim 13, wherein the acid comprises one or more of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, trifluoroacetic acid, p-

toluenesulphonic acid, maleic acid, acetic acid and succinic acid.

- [15] The process of claim 12, wherein the base comprises an inorganic or organic base.
- [16] The process of claim 15, wherein the base comprises one or more of an alkali and alkaline earth metal hydroxide or alkoxide; ammonia; triethylamine, dicyclohexylamine, diisopropylamine, or mixtures thereof.
- [17] The process of claim 12, further comprising isolating the acid addition salt of cefdinir.
- [18] The process of claim 12, wherein the crystalline cefdinir is the hydrated form of cefdinir.
- [19] The process of claim 12, wherein the crystalline cefdinir is either or both of Form B or Form C of cefdinir.
- [20] A process for the preparation of crystalline cefdinir, the process comprising:
(a) acidifying a base addition salt of cefdinir at a temperature of 10°C or less; and
(b) isolating the crystalline cefdinir.
- [21] The process of claim 20, wherein the acid comprises one or more of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, trifluoroacetic acid, p-toluenesulphonic acid, maleic acid, acetic acid and succinic acid.
- [22] A process for the preparation of crystalline Form B or Form C of cefdinir, the process comprising:
(a) dissolving cefdinir or salt thereof in water at pH of about 5.5 to 8;
(b) optionally adding an organic solvent in the solution of step a);
(c) adjusting the pH of the solution obtained in step a) or b) to between 1.0 and 3.0 with an acid; and
(d) isolating crystalline Form B of cefdinir from the reaction mixture obtained thereof.
- [23] The process according to claim 22, wherein the organic solvent is added in step b).
- [24] The process according to claim 23, wherein the organic solvent used in step b) is water miscible to get Form B.
- [25] The process according to claim 24, wherein the organic solvent used in step b) comprises one or more of methanol, ethanol, isopropanol, n-propanol, acetone, acetonitrile, tetrahydrofuran, 1,4-dioxane or mixtures thereof.
- [26] The process according to claim 22, wherein the organic solvent used in step b) comprises one or more of alkanols, esters, ketones, acetonitrile, chlorinated hydrocarbons, tetrahydrofuran, N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, 1,4-dioxane and dimethylsulphoxide or mixtures thereof to

get Form C.

- [27] A pharmaceutical composition comprising a therapeutically effective amount of one or both of the crystalline Form B and the crystalline Form C of cefdinir, and one or more pharmaceutically acceptable carriers, excipients or diluents.
- [28] The pharmaceutical composition according to claim 27, wherein the cefdinir comprises either substantially pure crystalline Form B cefdinir or substantially pure crystalline Form C cefdinir.
- [29] A method of treating bacterial infections in a warm-blooded animal, the method comprising providing a pharmaceutical composition to the warm-blooded animal, the pharmaceutical composition comprising one or both of the crystalline Form B and the crystalline Form C of cefdinir.
- [30] The method according to claim 29, wherein the cefdinir comprises either substantially pure crystalline Form B cefdinir or substantially pure crystalline Form C cefdinir.




FIGURE 2.




FIGURE 3





FIGURE 4



Paper Copy (NOT for submission)

FIGURE 5




FIGURE 6




FIGURE 7





FIGURE 8



INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB2005/052691

A. CLASSIFICATION OF SUBJECT MATTER
C07D501/00 A61K31/546 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2004/056835 A1 (ANTIBIOTICOS S.P.A., ITALY) 8 July 2004 (2004-07-08) cited in the application claims 8,9; example 5	1-5, 12-17, 19-25, 27-30
Y	US 2003/204082 A1 (MANCA, ANTONIO ET AL) 30 October 2003 (2003-10-30) cited in the application claim 2; example 1	1-5, 12-17, 19-25, 27-30
X	US 6 350 869 B1 (STURM, HUBERT ET AL) 26 February 2002 (2002-02-26) cited in the application example 3 (col.4, l.12-21); claim 1	1-5, 12-17, 19-25, 27-30
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 November 2005

Date of mailing of the international search report

27.12.05

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Schuemacher, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/182005/052691

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 093 814 A (LEE, GWAN SUN ET AL) 25 July 2000 (2000-07-25) examples 6-8 -----	1-5, 12-17, 19-25, 27-30
Y	US 4 935 507 A (TAKAYA ET AL) 19 June 1990 (1990-06-19) cited in the application claim 1; examples 1,4,5 -----	1-5, 12-17, 19-25, 27-30
X	WO 2004/016623 A1 (SANDOZ G.M.B.H., AUSTRIA) 26 February 2004 (2004-02-26) cited in the application example 1 -----	1-5, 12-17, 19-25, 27-30
P,X	WO 2004/104010 A1 (RANBAXY LABORATORIES LIMITED, INDIA) 2 December 2004 (2004-12-02) claim 14; examples 1,2 -----	1-30
P,X	US 2004/242556 A1 (DANDALA, RAMESH ET AL) 2 December 2004 (2004-12-02) cited in the application tables 1,2 -----	1-30
E	WO 2005/090360 A1 (ORCHID CHEMICALS & PHARMACEUTICALS LIMITED, INDIA) 29 September 2005 (2005-09-29) claims 1-3; examples 5-14 -----	1-30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2005/052691

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 29 and 30 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see annex

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-5,12-17(part.),19-25 (part.), 27-30(part.)

crystalline form B of cefdinir, the process for its preparation (claims 12-17 and 19-25, as far as they are related to the preparation of form B), the pharmaceutical composition comprising the crystalline form B and its use to treat bacterial infections.

2. claims: 6, 12-18(part.)

A hydrated crystalline form of cefdinir and the process for its preparation (claims 12-18, as far as they are related to the preparation of the hydrated crystalline form).

3. claims: 7-11, 12-17(part.), 19-23 (part.), 26,27-30(part.)

crystalline Form C of cefdinir, the process for its preparation (claims 12-17, 19-23 and 26, as far as they are related to the preparation of form C), the pharmaceutical composition comprising the crystalline form C and its use to treat bacterial infections.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB2005/052691

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004056835 A1	08-07-2004	AU 2003293746 A1 EP 1572699 A1	14-07-2004 14-09-2005
US 2003204082 A1	30-10-2003	CA 2424501 A1 JP 2004035543 A US 2005209451 A1	29-10-2003 05-02-2004 22-09-2005
US 6350869 B1	26-02-2002	AT 405283 B AT 57097 A AU 731413 B2 AU 7428898 A BR 9809745 A CA 2283718 A1 CN 1251590 A DE 69816056 D1 DE 69816056 T2 WO 9845299 A1 EP 0973779 A1 HU 0002987 A2 ID 22536 A JP 3421354 B2 JP 2000514833 T NO 994466 A PL 335620 A1 SK 134399 A3 TR 9902406 T2	25-06-1999 15-11-1998 29-03-2001 30-10-1998 20-06-2000 15-10-1998 26-04-2000 07-08-2003 15-04-2004 15-10-1998 26-01-2000 28-02-2001 04-11-1999 30-06-2003 07-11-2000 15-09-1999 08-05-2000 16-05-2000 21-02-2000
US 6093814 A	25-07-2000	AT 218572 T DE 69621649 D1 DE 69621649 T2 DK 874853 T3 EP 0874853 A1 ES 2175167 T3 JP 2000502700 T WO 9724358 A1 PT 874853 T	15-06-2002 11-07-2002 19-09-2002 23-09-2002 04-11-1998 16-11-2002 07-03-2000 10-07-1997 30-09-2002
US 4935507 A	19-06-1990	AT 123221 T AU 617347 B2 CA 1297096 C DE 3853901 D1 DE 3853901 T2 EP 0304019 A2 ES 2072856 T3 HK 18496 A IE 67348 B1 KR 9708126 B1 ZA 8805709 A	15-06-1995 28-11-1991 10-03-1992 06-07-1995 12-10-1995 22-02-1989 01-08-1995 09-02-1996 20-03-1996 21-05-1997 26-04-1989
WO 2004016623 A1	26-02-2004	AU 2003255424 A1 EP 1554289 A1	03-03-2004 20-07-2005
WO 2004104010 A1	02-12-2004	NONE	
US 2004242556 A1	02-12-2004	NONE	
WO 2005090360 A1	29-09-2005	NONE	